

loading of foramsulfuron and iodosulfuron-methyl from runoff water and sediment. Runoff of this product will be greatly reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

**GROUNDWATER ADVISORY**

This product has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the same area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**PRODUCT INFORMATION**

**PREVIOUS PRODUCT USES**

Derigo® Herbicide may be used as a foliar spray to control annual and perennial broadleaf weeds and grasses in unimproved warm season grasses including bermudagrass, centipedegrass, and zoysiagrass, and for use in bare ground sites. Unimproved grasses are grasses found in settings that are not well-maintained and DOES NOT include settings (e.g., sports fields, home lawns, golf courses) where the grass is maintained at a higher level through management practices including frequent mowing at short heights, and fertility. Derigo Herbicide can also be used to suppress seedheads and vegetative growth of bahiagrass\*\* in order to reduce mowing requirements.

Derigo Herbicide controls weeds after they have germinated (postemergence) and will provide short-term residual control of some listed annual broadleaf and grassy weeds.

Derigo Herbicide may be used to manage weeds on private, public and military non-crop lands (e.g. pipelines, railroads, utility and highway rights-of-way), airports, parks, natural areas, restoration sites, municipal sites (e.g., waste

water treatment facilities and prisons facilities), commercial and industrial sites (e.g. oil refineries, distribution centers, and chemical plants).

\*\*Not for use in California.

**SYMPTOMS**

Weed growth ceases within hours after a postemergence application. Symptoms progress from yellowing or red/purpling to eventual plant death within 1 to 4 weeks after application depending on the sensitivity of the weed and environmental conditions. Weed control is more rapid when average air temperatures are 65 degrees or greater, when soil moisture is adequate for weed growth, and when weeds are not under environmental stress.

**WARM SEASON GRASS TOLERANCE**

Derigo Herbicide can be used on the following types of low maintenance grasses: bermudagrass (*Cynodon dactylon*), centipedegrass (*Eriochloa ophioides*), and zoysiagrass (*Zoysia species*). All of these species show good tolerance to this product, however, some temporary discoloration of certain warm-season grasses may occur to turf under stress from drought, disease, extreme cold or hot weather.

Derigo Herbicide will suppress bahiagrass\*\* (*Paspalum notatum*) growth and seedhead development. Bahiagrass\*\* may be temporarily discolored following application depending on bahiagrass\*\* cultivars, herbicide rate, surfactant used, application timing, and environmental factors.

Other warm season grasses and their cultivars may be tolerant to this product, however tolerance testing needs to be done prior to use.

**DO NOT** use this product on cool-season grasses, including tall fescue, fine fescue, Kentucky bluegrass, perennial ryegrass, etc. unless control of these species is desired.

\*\* Not for use in California.

**MODE OF ACTION**

Derigo Herbicide contains three active ingredients: thienacarbazon-methyl, iodosulfuron-methyl-sodium, and foramsulfuron. These active ingredients inhibit acetolactate synthase (ALS), an enzyme responsible for the synthesis of amino

acids that are essential for plant growth. Some weed species have naturally occurring biotypes that are resistant to ALS-inhibiting herbicides.

**RESISTANCE MANAGEMENT**

Derigo Herbicide contains the active ingredients foramsulfuron, iodosulfuron-methyl, and thienacarbazon-methyl which are Group 2 Herbicides based on the mode of action classification system of the Weed Science Society of America. When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. Adequate control of these resistant weed biotypes cannot be expected.

Follow the best management practices listed below to delay the development of herbicide resistant weeds.

- Fields should be scouted prior to application to identify the weed species present and their growth stage to determine if the intended application will be effective. Fields should be scouted after application to verify that the treatment was effective.
- Identify weeds present in the field through scouting and field history and understand their biology. The weed-control program should consider all of the weeds present.
- Suspected herbicide-resistant weeds may be identified by these indicators:
  - o Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
  - o A spreading patch of non-controlled plants of a particular weed species; and
  - o Surviving plants mixed with controlled individuals of the same species.
- Contact your local sales representative, crop advisor, or extension agent to find out if suspected resistant weeds to this MOA have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective mechanisms of actions for each target weed.

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

**For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-331-2867**

FIRST AID	
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>Immediately call a poison control center or doctor for treatment advice.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>

**For MEDICAL Emergencies Call 24 Hours A Day 1-800-331-2867. Have the product container or label with you when calling a poison control center or doctor or going for treatment.**

**PRECAUTIONARY STATEMENTS**  
**HAZARD TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION**  
 Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or on clothing. Wash thoroughly with soap and

FORAMSULFURON & IODOSULFURON-METHYL & THIENCARBAZONE-METHYL		GROUP 2	HERBICIDE
---	--	---------	-----------

Used to manage weeds in Unimproved Warm-Season Grass Areas, Non-irrigation ditch banks, Pipelines, Highways, Airports, Railroads, Utility Rights-of-Way, Parks, Natural areas, Military Installations, Restoration sites, Municipal sites, Manufacturing sites, Sewage disposal sites, Commercial sites, and Industrial sites. Also used for Seedhead and Vegetative Growth Regulation of bahiagrass\*\*.

**ACTIVE INGREDIENTS:**  
 Foramsulfuron (CAS Number 173159-57-4)..... 24.0%  
 Iodosulfuron-methyl (CAS Number 146650-36-7)..... 2.4%  
 Thienacarbazon-methyl (CAS Number 317815-83-1)..... 10.0%  
**OTHER INGREDIENTS:**..... 65.6%  
**TOTAL:**..... 100.0%

Derigo® Herbicide is formulated as a 36.4% water dispersible granule

EPA Reg. No. 432-1533 EPA Est. No. 000264-DEU-001  
 See Back Panel for First Aid Instructions and Booklet for Complete Precautionary Statements and Directions for Use.

**STOP - READ THE LABEL BEFORE USE**  
**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

Produced for:  
 Bayer Environmental Science  
 Division of Bayer CropScience, Inc.  
 5000 Crofton Green Way, Suite 400  
 Cary, NC 27513  
 Product of Germany

Net Contents  
 60 Ounces (1.76kg)  
 81785031  
 84127191C  
 191204AV1



water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

- All mixers, loaders, applicators and other handlers must wear:
- Long-sleeved shirt and long pants
  - Shoes plus socks
  - Waterproof gloves

**USER SAFETY RECOMMENDATIONS**

- Users should:**
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
  - Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This product is toxic to non-target plants. Non-target plants may be adversely affected if the product is allowed to drift from the areas of application. Avoid spray drift from treated area. Do not apply when conditions favor drift from treated areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate. Do not drain or rinse equipment near desirable vegetation. Refer to the Spray Drift Management section of this label for additional information.

**SURFACE WATER ADVISORY**

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for weeks or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential

continued

3. Apply spray mixtures of this product within 5 days of mixing to avoid product degradation.
- APPLICATION METHODS**
- Uniform, thorough spray coverage of the targeted weeds with properly calibrated spray equipment is important to achieve consistent weed control. Derigo Herbicide may be applied by broadcast or spot applications.
- Broadcast Application**
- Apply Derigo Herbicide at a rate of 3 to 6 ounces per acre as a broadcast spray to control the weeds listed in the WEEDS CONTROLLED Section of the label. **DO NOT** exceed the maximum amount of 6 ounces per acre of this product per year applied as a broadcast spray. For broadcast applications, use a minimum of 10 gallons of water per acre. A higher spray volume may be necessary to obtain adequate spray coverage of targeted weeds for control in dense weed populations.
- Spot Application/ Directed Spray**
- Apply with a hand held or high volume application sprayer. Apply a spray solution consisting of 3-6 ounces product per 25 to 100 gallons as directed spray to targeted weeds until wet. For spot treatment, **DO NOT** treat more than 1000 sq. ft. per acre of treated area.
- Non-irrigation Ditch Banks**
- It is permissible to treat non-irrigation ditch banks and transitional areas between upland and lowland sites. Derigo Herbicide can be used to the water's edge. **DO NOT** apply directly to water and take precautions to minimize spray drift into water. Refer to spray drift management section of this label for more information.
- USE OF SPRAY ADJUVANTS**
- Derigo Herbicide is a water dispersible granule (WDG) that requires the use of a surfactant for maximum weed control. Use the spray adjuvant(s) as specified in the WEED CONTROL section of this label or follow the directions below.
- Use a non-ionic surfactant (NIS) at 0.25 to 0.5% v/v of the spray solution that is at least 80% active material. **DO NOT** exceed 1 quart of NIS per acre

- as injury to desirable grass species may occur.
  - Use of a spray adjuvant containing an organosilicone surfactant is NOT advised.
  - For difficult-to-control weeds and perennial grasses, use 0.5 to 1% v/v of a methylated seed oil (MSO) containing at least 80% methylated seed oil.
  - The addition of ammonium sulfate (AMS) or urea ammonium nitrate (UAN) has been shown to improve control of some difficult-to-control weeds. Use a spray grade AMS (1.5 to 3 lb/A) for areas of high relative humidity or use UAN (1.5-2 qt/A) in areas of low relative humidity.
  - Application of Derigo Herbicide with a spray adjuvant or nitrogen-containing fertilizers when temperatures are above 90 degrees or when desirable grasses are under stress may cause injury.
- MIXING, COMPATIBILITY, AND CLEANUP**
- Spray Solution pH**
- It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.
- The efficacy of this product may be affected by the pH of the spray solution. A pH near 7.0 is ideal. If the pH is <6 and if product spray solution is not to be used within 24 hours, add a suitable buffer.
- Mixing Instructions**
- This product must be applied with clean and properly calibrated equipment. Prior to adding this product, ensure that the spray tank, filters, and nozzles have been thoroughly cleaned. Prepare only as much spray mixture as needed for application on the same day.
- Fill spray tank with 25% to 50% of the required volume of water, and begin agitation prior to the addition of this product.
  - Before filling or adding any additional products, ensure full dispersion of this product.

- MANDATORY SPRAY DRIFT (continued)**
- For all applications, applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
  - Do not apply when wind speeds exceed 10 miles per hour at the application site.
  - Do not apply during temperature inversions.
  - Bottom-Less Ground Applications:
    - Applicators are required to use a Medium or coarser droplet size (ASABE S572.1) for all applications.
    - Do not apply when wind speeds exceed 10 miles per hour at the application site.
    - Do not apply during temperature inversions.

- SPRAY DRIFT ADVISORIES**
- Bottom-Less Ground Applications:
- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.
- Handheld Technology Applications:
- Take precautions to minimize spray drift.
- THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS. IMPORTANCE OF DROPLET SIZE**
- An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made in properly or under unfavorable environmental conditions.
- Controlling Droplet Size - Ground Boom**
- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.

- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
  - Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.
- Controlling Droplet Size - Aircraft**
- Adjust Nozzles - Follow nozzle manufacturer's recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.
  - BOOM HEIGHT - Ground Boom Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.
- RELEASE HEIGHT - Aircraft**
- Higher release heights increase the potential for spray drift. When applying aerially to crops, do not release spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.
- SHIELDED SPRAYERS**
- Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.
- TEMPERATURE AND HUMIDITY**
- When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.
- TEMPERATURE INVERSIONS**
- Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

- WIND**
- Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.
- NON-TARGET ORGANISM ADVISORY**
- This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated area. Protect the forage and habitat of non-target organisms by minimizing spray drift. For further guidance and instructions on how to minimize spray drift, refer to the Spray Drift Management section of this label.
- WINDBLOWN SOIL PARTICLES ADVISORY**
- Derigo Herbicide has the potential to move off-site due to wind erosion. Soils that are subject to wind erosion usually have a high silt and/or fine to very fine sand fractions and low organic matter content. Other factors which can affect the movement of windblown soil include the intensity and direction of prevailing winds, vegetative cover, site slope, rainfall, and drainage patterns. Avoid applying Derigo Herbicide if prevailing local conditions may be expected to result in off-site movement.
- RESTRICTIONS**
- DO NOT** apply more than a total of 6 ounces (0.090 lb foramsulfuron; 0.009 lb iodossulfuron-methyl; 0.0375 lb thiencarbazone-methyl) product per acre per year (365 days).
  - DO NOT** apply more than 6 ounces (0.090 lb foramsulfuron; 0.009 lb iodossulfuron-methyl; 0.0375 lb thiencarbazone-methyl) of product per acre in a single application.
  - For Bahiagrass growth regulation, **DO NOT** apply more than a total of 3 ounces (0.045 lb foramsulfuron; 0.0045 lb iodossulfuron-methyl; 0.0188 lb thiencarbazone-methyl) product per acre per year (365 days).
  - DO NOT** make more than two applications of Derigo Herbicide to all use sites per year when using reduced application rates. Allow 4 to 6 weeks between applications.
  - DO NOT** apply this product by air or through any type of irrigation system.

- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
  - Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.
- Controlling Droplet Size - Aircraft**
- Adjust Nozzles - Follow nozzle manufacturer's recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.
  - BOOM HEIGHT - Ground Boom Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.
- RELEASE HEIGHT - Aircraft**
- Higher release heights increase the potential for spray drift. When applying aerially to crops, do not release spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.
- SHIELDED SPRAYERS**
- Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.
- TEMPERATURE AND HUMIDITY**
- When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.
- TEMPERATURE INVERSIONS**
- Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

6. **DO NOT** use on residential turf, sod farms, golf courses, athletic fields or other improved turf settings.
10. **DO NOT** use Derigo Herbicide for weed control in pastures and hayfields.
11. **DO NOT** use this product on cool-season grass types, including tall fescue, fine fescue, Kentucky bluegrass, perennial ryegrass unless control of these species is desired.
12. **DO NOT** mow immediately after application or before spray has dried or weed control may be reduced. After treatment, do not transfer clippings to non-target areas.
- Amount of active ingredients for select Derigo Herbicide rates.
- | Derigo Herbicide Rate ounces/A | Foramsulfuron lb ai/A | Iodossulfuron-methyl lb ai/A | Thiencarbazone-methyl lb ai/A |
|--------------------------------|-----------------------|------------------------------|-------------------------------|
| 1.5                            | 0.023                 | 0.0023                       | 0.0094                        |
| 3                              | 0.045                 | 0.0045                       | 0.0188                        |
| 6                              | 0.090                 | 0.0090                       | 0.0375                        |
- PRECAUTIONS**
- Rainfall within 2 hours of spray drying may result in reduced weed control and may necessitate retreatment.
  - Weeds need to be actively growing when the herbicide application is made. Mature, hardened-off weeds may not be controlled. Weed control may be reduced if application is made when weeds are dust covered or in the presence of heavy dew, fog, and mist/rain or when weeds are under stress due to drought.

3. If this product is applied in a tank mixture with other products, add this product to the spray tank first and ensure it is thoroughly dispersed before adding other products.
4. Continue to fill the spray tank with water to the desired volume and agitate while adding spray adjuvants or nitrogen fertilizers.
- Compatibility**
- This product is to be tank-mixed with other products. It is advised that a compatibility test be done prior to mixing. To test for compatibility, use a small container and mix a small amount (0.5 to 1 qt) of spray, combining all ingredients in the same ratio as the anticipated use. If any indications of physical incompatibility develop (precipitation, settling, changes in color) do not use this mixture for spraying. Indications of incompatibility may occur within 5-15 minutes after mixing. Read and follow the label of each tank mix product used for precautionary statements, directions for use, geographic and other restrictions.
- Tank Mixture Procedure**
- Drain the tank completely, then wash out tank, boom, and hoses with clean water. Drain again.
  - Fill the tank full with clean water and add ammonia (i.e. 3% domestic ammonia solution) at a dilution rate of 1% (i.e. 1 gallon of domestic ammonia for every 100 gallons of rinse). Completely fill the tank with water. Agitate-rinse and flush through boom and hoses. Leave agitation on for 10 minutes. Drain tank completely.
  - Repeat Step 2.
  - Remove nozzles and screens and soak them in a 1% ammonia solution. Inspect nozzles and screens and remove visible residues.
  - Flush tank, boom, and hoses with clean water.
  - Inspect tank for visible residues. If present, repeat Step 2.
- WEED CONTROL INFORMATION**
- This product may be used to control a variety of broadleaf weeds and grasses in warm season perennial grasses listed on this label. Apply this product to

- susceptible weeds as listed in the Weed Control section of this label. For certain perennial and difficult to control annual weeds, a repeat application may be needed 4 to 6 weeks later if regrowth is observed. Total amount of product applied in a calendar year (365 days) must not exceed 6 ounces of product per acre.
- VASEYGRASS, DALLISGRASS, and JOHNSONGRASS CONTROL**
- For best results, apply this product in combination with MSU at 0.5 to 1% v/v and apply as a broadcast, spot application, or directed spray. Make a second application if regrowth is observed 30 to 60 days later. Applications made in the fall generally provide better control of dallisgrass than at other times of the year.
- BAHAGRASS\*\* GROWTH REGULATION**
- Derigo Herbicide may be applied as a broadcast spray to bahiagrass at a rate of 1.5 to 3 ounces per acre to suppress tiller and seedhead growth. Higher use rates, including those used for weed control (greater than 3 ounces per acre), may cause unacceptable bahiagrass discoloration. For optimum results, apply after bahiagrass green-up and avoid applications when bahiagrass is under visible signs of stress due to drought or other environmental factors. Length of growth regulation will depend on use rate, bahiagrass cultivar, and environmental conditions.
- \*\*Not for use in California.
- TANK-MIXES**
- Derigo Herbicide may be tank-mixed with (but not limited to) the following herbicides to provide extended residual control or improved postemergence weed control: Esplanade® 200SC Herbicide (EPA Reg. No. 432-1516 containing indaziflam), aminopyralid, aminocyclopyrachlor, glyphosate, triclopyr.
- When using Derigo Herbicide in combination with other herbicides, follow the precautions, restrictions and directions of both labels. When using new tank mixtures with Derigo Herbicide, test physical and biological compatibility prior to use.**

7. If this product is applied in a tank mixture with other products, add this product to the spray tank first and ensure it is thoroughly dispersed before adding other products.
8. Continue to fill the spray tank with water to the desired volume and agitate while adding spray adjuvants or nitrogen fertilizers.
- Compatibility**
- This product is to be tank-mixed with other products. It is advised that a compatibility test be done prior to mixing. To test for compatibility, use a small container and mix a small amount (0.5 to 1 qt) of spray, combining all ingredients in the same ratio as the anticipated use. If any indications of physical incompatibility develop (precipitation, settling, changes in color) do not use this mixture for spraying. Indications of incompatibility may occur within 5-15 minutes after mixing. Read and follow the label of each tank mix product used for precautionary statements, directions for use, geographic and other restrictions.
- Tank Mixture Procedure**
- Drain the tank completely, then wash out tank, boom, and hoses with clean water. Drain again.
  - Fill the tank full with clean water and add ammonia (i.e. 3% domestic ammonia solution) at a dilution rate of 1% (i.e. 1 gallon of domestic ammonia for every 100 gallons of rinse). Completely fill the tank with water. Agitate-rinse and flush through boom and hoses. Leave agitation on for 10 minutes. Drain tank completely.
  - Repeat Step 2.
  - Remove nozzles and screens and soak them in a 1% ammonia solution. Inspect nozzles and screens and remove visible residues.
  - Flush tank, boom, and hoses with clean water.
  - Inspect tank for visible residues. If present, repeat Step 2.
- WEED CONTROL INFORMATION**
- This product may be used to control a variety of broadleaf weeds and grasses in warm season perennial grasses listed on this label. Apply this product to

**Broadleaf Weeds Controlled** (continued)

Common Name	Scientific Name
Prickly sida*	<i>Sida spinosa</i>
Prostrate knotweed	<i>Polygonum aviculare</i>
Purple cloverweed	<i>Chamaesyce maculata</i>
Purple deadnettle	<i>Griffithalia purpureum</i>
Rabbitfoot clover	<i>Lamium purpureum</i>
Red sorrel	<i>Rumex acetosella</i>
Redroot pigweed	<i>Amaranth retrofractus</i>
Russian thistle	<i>Salsola tragus</i>
Shepherd's purse	<i>Capsella bursa-pastoris</i>
Sicklepod	<i>Senna obtusifolia</i>
Slender aster	<i>Aster gracilis</i>
Southern brassbuttons	<i>Crotalaria australis</i>
Spiny sowthistle	<i>Sonchus asper</i>
Spreading horsetweed	<i>Callitriche vialis</i>
Swinecress	<i>Coronopus didymus</i>
Tansy mustard	<i>Descurainia pinnata</i>
Turnipweed	<i>Rapistrum rugosum</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Venus looking-glass	<i>Triodanis perfoliata</i>
Virginia buttonweed*	<i>Diodia virginiana</i>

continued

**Broadleaf Weeds Controlled** (continued)

Common Name	Scientific Name
Virginia dwarf dandelion	<i>Krigia virginica</i>
Western ragweed	<i>Ambrosia psilostachya</i>
White clover	<i>Trifolium repens</i>
White mustard	<i>Brassica alba</i>
White sweet clover	<i>Melilotus alba</i>
Whiteleaf sage	<i>Salvia leucophylla</i>
Wild buckwheat	<i>Polygonum convolvulus</i>
Wild carrot	<i>Daucus carota</i>
Wild garlic	<i>Allium vineale</i>
Wild lettuce	<i>Lactuca canadensis</i>
Wild mustard	<i>Brassica kaber</i>
Wild onion	<i>Allium canadense</i>
Wild parsley	<i>Lomatium tomentosum</i>
Wild radish	<i>Raphanus raphanistrum</i>
Yellow rocket	<i>Barbarea vulgaris</i>
Yellow woodsorrel, Oxalis*	<i>Oxalis stricta</i>

\*\*Not for use in California.

**USE RATES FOR WEED CONTROL**

Apply this product at a rate of 3 to 6 ounces per acre as a broadcast spray or at a rate of 3 to 6 ounces per 25 to 100 gallons as a directed spray to control the weeds listed below. Some weed species and more mature weed growth stage may require repeat applications and/or use of the higher use rate on this product label even under ideal conditions for application.

**Broadleaf Weeds Controlled**

Common Name	Scientific Name
American burweed, Fireweed	<i>Erechtites hieracifolia</i>
Annual lespedeza	<i>Lespedeza striata</i>
Asiatic hawkbeard	<i>Youngia japonica</i>
Bircheye pearwort	<i>Sagina procumbens</i>
Black medic, hop medic	<i>Medicago lupulina</i>
Black nightshade	<i>Solanum nigrum</i>
Blackseed plantain	<i>Plantago rugelii</i>
Blue mustard	<i>Chorospora tenella</i>
Broadleaf plantain	<i>Plantago aristata</i>
Broadleaf plantain, common plantain	<i>Plantago major</i>
Buckhorn plantain, narrowleaf plantain	<i>Plantago lanceolata</i>
Burcucumber	<i>Sisyrinchium angustatum</i>
Burdock	<i>Arctium spp.</i>
Buttercup	<i>Ranunculus spp.</i>
California burclover	<i>Medicago polymorpha</i>
Chamomile, scentless	<i>Matricaria inodora</i>

continued

**Grassy Weeds and Sedges Controlled**

Common Name	Scientific Name
Annual bluegrass	<i>Poa annua</i>
Annual ryegrass	<i>Lolium multiflorum</i>
Barnyardgrass	<i>Echinochloa crusgalli</i>
Broadleaf signalgrass	<i>Urochloa platyphylla</i>
Browntop millet	<i>Brachiaria ramosa</i>
Carpetgrass	<i>Axonopus affinis</i>
Common millet, proso millet	<i>Panicum miliaceum</i>
Giant foxtail	<i>Setaria faberi</i>
Goosegrass	<i>Elymus indica</i>
Green foxtail	<i>Setaria viridis</i>
Gopher tail grass	<i>Eragrostis ciliaris</i>
Green kyllinga	<i>Kyllinga brevifolia</i>
Dallisgrass*	<i>Paspalum dilatatum</i>
Doveweed	<i>Murdannia nudiflora</i>
Fall panicum	<i>Panicum dichotomiflorum</i>
Field sandbur	<i>Cenchrus incertus</i>
Johnsongrass*	<i>Sorghum halepense</i>
Large crabgrass*	<i>Digitaria sanguinalis</i>
Quackgrass	<i>Agropyron repens</i>
Red fescue	<i>Festuca rubra</i>
Rental fescue	<i>Vulpia myuros</i>

continued

**Grassy Weeds and Sedges Controlled** (continued)

Common Name	Scientific Name
Rescuegrass*	<i>Bromus catharticus</i>
Shattercane	<i>Sorghum bicolor</i>
Stinkgrass	<i>Eragrostis cilianensis</i>
Switchgrass	<i>Panicum virgatum</i>
Tall fescue	<i>Festuca arundinacea</i>
Texas panicum	<i>Panicum texanum</i>
Thin paspalum, bull paspalum*	<i>Paspalum setaceum</i>
Vaseygrass*	<i>Paspalum urvillei</i>
Wild oat	<i>Avena fatua</i>
Yellow foxtail	<i>Setaria luberosa</i>

\* Weeds may require a second application of this product for acceptable control. The degree of weed control varies with rate, weed size or stage of growth, and environmental conditions before and following treatment. If weeds are showing signs of recovery, make a second application 4 to 6 weeks after the first. **DO NOT** exceed 6 ounces of product per acre per year (365 days) for all applications.

**Broadleaf Weeds Controlled** (continued)

Common Name	Scientific Name
Canada thistle**	<i>Cirsium arvense</i>
Canada trefoil	<i>Linaris canadensis</i>
Carolina dichondra, Dichondra*	<i>Dichondra carolinensis</i>
Carolina false-dandelion	<i>Pyrrhopappus carolinianus</i>
Carolina geranium, wild geranium*	<i>Geranium carolinianum</i>
Carpetweed	<i>Mollugo verticillata</i>
Cats ear dandelion	<i>Hypochaeris radicata</i>
Chamberbitter	<i>Phyllanthus urinaria</i>
Common chickweed	<i>Stellaria media</i>
Common lambsquarters*	<i>Chenopodium album</i>
Common purslane*	<i>Portulaca oleracea</i>
Common ragweed	<i>Ambrosia artemisiifolia</i>
Common sunflower	<i>Helianthus annuus</i>
Common vetch	<i>Vicia sativa</i>
Common waterhemp	<i>Amaranthus rudis</i>
Common speedwell	<i>Veronica anensis</i>
Creeping beggarweed	<i>Desmodium illinoense</i>
Creeping speedwell	<i>Veronica filiformis</i>
Curly dock	<i>Rumex crispus</i>
Cutleaf evening primrose	<i>Oenothera laciniata</i>
Dandelion, common	<i>Taraxacum officinale</i>

continued

**Broadleaf Weeds Controlled** (continued)

Common Name	Scientific Name
Hempnettle	<i>Galeopsis spp.</i>
Henbit	<i>Lamium amplexicaule</i>
Hop clover	<i>Trifolium spp.</i>
Horse-nettle	<i>Solanum carolinense</i>
Horse purslane	<i>Trianthema portulacastrum</i>
Horseweed (marshall)**	<i>Conyza canadensis</i>
lytle morningglory	<i>Ipomoea hederacea</i>
Khakiweed*	<i>Alternanthera caracasana</i>
Knael	<i>Sceleranthus annuus</i>
Lady's Mantle	<i>Alchemilla mollis</i>
Lawn burweed, spurweed	<i>Sonchus oleraceus</i>
Mouse-ear chickweed	<i>Cerastium glomeratum</i>
Oxeye daisy	<i>Leucanthemum vulgare</i>
Palseed plantain	<i>Plantago virginica</i>
Palmer amaranth	<i>Amaranth palmeri</i>
Parsley plect	<i>Aphanes microcarpa</i>
Pennsylvania smartweed**	<i>Polygonum pensylvanicum</i>
Pitted morningglory	<i>Ipomoea lacunosa</i>
Poison hemlock	<i>Conium maculatum</i>
Pokeweed, common	<i>Phytolacca americana</i>
Poolrue*	<i>Dicella tetras</i>

continued

**Broadleaf Weeds Controlled** (continued)

Common Name	Scientific Name
Dogfennel	<i>Eupatorium capillifolium</i>
Dollarweed, Pennywort*	<i>Hydrocotyle spp.</i>
Eastern black nightshade	<i>Solanum ptychanthum</i>
Entireleaf morningglory	<i>Ipomoea hederacea var. integrifolia</i>
Facelis, trampweed	<i>Facelis retusa</i>
Field madder	<i>Sherardia arvensis</i>
Field pansy, Johnny jump-up*	<i>Viola rufinesquibicolor</i>
Field pepperweed	<i>Lepidium campestre</i>
Field pennycress	<i>Thlaspi arvense</i>
Field violet, wild pansy	<i>Viola arvensis</i>
Field bindweed	<i>Erigeron spp.</i>
Florida betony	<i>Stachys floridana</i>
Florida pusley	<i>Richardia scabra</i>
Giant ragweed	<i>Ambrosia trifida</i>
Ground ivy	<i>Glechoma hederacea</i>
Harry bittercress	<i>Cardamine hirsuta</i>
Harry nightshade	<i>Solanum villosum</i>
Heartwing sorrel	<i>Rumex hastatulus</i>
Heath aster**	<i>Aster ericoides</i>
Hemp sesbania	<i>Sesbania exaltata</i>

continued

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

### PESTICIDE STORAGE

Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

### PESTICIDE DISPOSAL

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

### CONTAINER HANDLING

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinse into application equipment or a mix tank or store rinse for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning, if burned, stay out of smoke.

## CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY AND LIABILITY

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, plant injury, other prop-

erty damage, as well as other unintended consequences may result because of factors beyond the control of Bayer CropScience LP. Those factors include, but are not limited to, weather conditions, presence of other materials or the manner of use or application. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

Bayer (reg'd), the Bayer Cross (reg'd), Esplanade® and Derigo® are registered trademarks of Bayer.

E-29 is a registered trademark of Nufarm.

12



## Derigo® HERBICIDE

### ACTIVE INGREDIENTS:

Foramsulfuron (CAS Number 173199-57-4).....24.0%

Iodosulfuron-methyl (CAS Number 144550-36-7).....2.4%

Thiencarbazone-methyl (CAS Number 317815-63-1).....10.0%

**OTHER INGREDIENTS:**.....33.6%

**TOTAL:**.....100.0%

Derigo® Herbicide is formulated as a 36.4% water dispersible granule

EPA Reg No. 432-1533 EPA Est. No. 000284-DEU-001

STOP - READ THE LABEL BEFORE USE

KEEP OUT OF REACH OF CHILDREN

**CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a

(If you do not understand the label, find someone to explain it to you in detail.)

See Panel for First Aid Instructions and Booklet for Complete

Precautionary Statements and Directions for Use.

Produced for:

Bayer Environmental Science

A Division of Bayer CropScience LP

5000 CentreGreen Way, Suite 400

Cary, NC 27513

81785031

8412719C.191204A11 Product of Germany

FORAMSULFURON & IODOSULFURON-METHYL & THIENCARBAZONE-METHYL

GROUP 2

HERBICIDE

For MEDICAL and TRANSPORTATION Emergencies ONLY Call

24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-800-331-2867

### FIRST AID

**If in eyes:**

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

**If on skin or clothing:**

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

**If swallowed:**

- Immediately call a poison control center or doctor for treatment advice.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Have person sip a glass of water if able to swallow.
- Do not give anything by mouth to an unconscious person.

For MEDICAL Emergencies Call 24 Hours A Day 1-800-334-7577. Have the product container with you when calling a poison control center or doctor or going for treatment.

Bayer